]•}Œ Œ š]vP v^µvŽšvoš Z hÇ
u}vP }oo P ^šµ vš•

^µuu ŒÇ }(šZ Z vš >]š Œ šµŒ

July 2011

This summary review was prepared for the BC Partners for Mental Health and Addictions Information to inform future planning and product development related to ongoing work supporting efforts to promote positive mental health on BC post-secondary campuses

Authors:

Tim Dyck is a Research Associate at the Centre for Addictions Research of BC, University of Victoria.

Dan Reist is the Assistant Director (Knowledge Exchange) at the Centre for Addictions Research of BC, University of Victoria.

resulting in intoxication) have become a growing concern in regard to young women (Kelly-Weeder, 2008; Grucza et al., 2009). Formal diagnostic designations of alcohol or other substance use disorders fall into two brackets: dependence and abuse (DSM-IV).

Alcohol dependence (or substance dependence) requires three or more of the following symptoms measured within the past year:

anorexia nervosa, a rate similar to the general population, to 46% in bulimia nervosa (Woodside & Staab, 2006; Bulik et al., 2004; Hasin et al., 2007).

These

unhealthy attitudes like desiring weight loss though maintaining a normal level (Adams- ï ‘••‘” ‡– al., 2006).Compared to their schoolmates who are not showing symptoms, those who are involved in disordered eating patterns suffer more harm related to use of other substances as well as in regard to alcohol (Dunn et al., 2002, 2009).

demonstration of such processes is required to substantiate a role in helping to precipitate or exacerbate such disorders (Wolfe & Maisto, 2000; Grilo, [illegible] & Marlatt, 2010; Ferriter & Ray, 2011).

Common predisposing personality characteristics

Some have contended that certain personality traits lend themselves to individuals falling prey to compulsions to eat and to indulge in substances (Ferriter & Ray, 2011). This claim, however, is questioned by others because the underlying premises (that eating disorders can be characterized as addictive behavioural patterns and that personality traits can be identified in individuals with eating disorders and substance use problems that make them vulnerable to these two addictions) have not been demonstrated.

Attention has been drawn to common markers such as increasing loss of behavioural control, preoccupation with the stuff being consumed, reliance on intake to cope with negative affect, and temporary emotional gratification followed by chronic harm. But these features do not characterize all eating disorders or all substance use problems. Even the notion of an addictive substance is dubious in regard to food and at least some drugs. [illegible] profile, and the bulimic bent to avoid food and control intake runs in conspicuous contrast to an attraction and lack of restraint associated with chronic drinking (Wolfe & Maisto, 2000).

Certain types or styles of personality (such as sensation-seeking, rebelliousness, low self esteem and depressed mood) have been identified with problematic patterns of substance use, but the notion of an addictive personality as a direct cause of substance use problems has been generally rejected (Chiauzzi & Liljegren, 1993). Just as there is an absence of uniformity in relationships between personality types and substance use problems, there is enough variety of (and even a striking contrast in) personality traits among those with disordered eating to undermine a conclusion of causal relationship. The strongest evidence supporting the role of personality in the relationship between eating disorders and substance use comes from studies that found that individuals with comorbid eating and substance use disorders are more likely to be diagnosed with borderline personality disorder and to report a history of self-destructive behaviour such as attempted suicide, promiscuity and reckless driving (Grilo et al., 2002; Sansone et al., 1994) compared to eating disordered individuals without a co-occurring substance use disorder. However, this may simply reflect the fact that the diagnostic criteria for borderline personality disorder include items that overlap with bulimic and substance use symptomatology.

Research indicating behavioural dysregulation is more predictive of substance use disorders, while avoidance and insecurity along with obsessional sensitive traits, figure more in those developing eating disorders, makes shared personality features less than satisfactory as an underlying explanation of acquisition. This is all the more reinforced when other factors like substance use history show themselves more successful in predicting later occurrence of difficulties (Wolfe & Maisto, 2000; Thompson-Brenner et al., 2008; Harrop & Marlatt, 2010; cf. Stewart & Brown, 2007a).

Comparable impact of a conducive family history

Another candidate to explain both eating and substance use problems is family history, whether genetic or experiential. Family history is widely acknowledged as a factor in regard to substance use disorders (and alcohol use disorders in particular), but the evidence in regard to eating

disorders is unclear (Wolfe & Maisto, 2000). The proposition that eating disorders and substance use disorders have a common genetic component has been challenged by results from a large-scale female twin study (Kendler et al., 1995). Moreover, cross-transmission, whereby eating disorder symptoms in one generation are associated with substance use disorders in the next and substance use difficulties for parents with eating difficulties in children, simply lacks evidence to confirm it as a vehicle ‘Ž ˆ ‡ ¬ ƒ ‹ • – ‘ á t r r r â ƒ ” ” ‘ ’ ¬ ƒ ” Ž ƒ – – á t r s r â … ˆ ä ” ‹ Ž ‘ ‡ – ƒ Ž ä á t 2000).

Certainly, family contexts characterized by chronic and traumatic parental alcohol-associated problems are liable to deleterious impact on personal development for children, but the current evidence does not demonstrate underlying familial mechanisms to link problematic substance use with the development of disordered eating. There remains a sizable prevalence of co-occurrence for individuals not subjected to such family history factors (Wolfe & Maisto, 2000; Harrop & Marlatt, 2010; cf. Stewart & Brown, 2007).

Common developmental

attestation that would widely establish such utilization as a primary motivation and proven means (Wolfe & Maisto, 2000).

Substances serving to compensate for self-starving

Food deprivation has been cited as a precipitator of substance use. Removal of food as a primary reinforcer would raise the value of alternative reinforcers such as alcohol or other drugs. Repeated experience of such hunger-stimulus satisfaction would consolidate their appeal. Here again, however, there is a lack of supporting evidence to corroborate this explanation, and ethical constraints limit experimental exploration of such a factor (Wolfe & Maisto, 2000; Harrop & Marlatt, 2010; … ˆ ä ” ‹ Ž ‘ ‡ – ƒ Ž ä á t r r t â ‹ • Š ƒ ¬ ï ƒ Ž Ž ‡ › á t r r r ä

Conclusion on etiology

The co-occurrence of difficulties with eating and substance use cannot yet be satisfactorily accounted for in terms of any particular type of contributing factor proving typically predominant (Harrop & Marlatt, 2010; Wolfe & Maisto, 2000; Grilo et al., 2002; Stewart & Brown, 2007a; Ferriter & Ray, 2001). A broader biopsychosocial framework, with attentiveness to a plurality of dynamics that can variously bear on persons and prompt the behaviours, is preferable for situating the phenomena. The frequent impact of trauma, especially in the form of sexual abuse, is not to be ignored. E ˜ ‡ • – Š ‘ — ‰ Š ‹ – ‡ ˜ ‹ † ‡ • – in a all cases, rates of physical and sexual abuse are significantly elevated in eating disordered populations, and where substance use disorders factor in as well, the abuse rates are doubled again (Harrop & Marlatt, 2010; Stewart & Brown, 2007a,b).

The diversity of substance use problems, as well as patterns of disordered eating, suggests the examination of more specific connections between certain drug use patterns and particular practices (bingeing, purging) will shed more light on pertinent factors than merely observing associations between undifferentiated use and established major types of eating disorders like anorexia nervosa or bulimia nervosa (Harrop & Marlatt, 2010; Dunn et al., 2009; Piran & Robinson, 2006a,b; Wolfe & Maisto, 2000; Corte & Stein, 2000; Wiederman & Pryor, 1996). One example of a more focused examination has detected both correlation and divergence between binge eating and heavy drinking. Both act as negative reinforcement, bringing relief to states of unpleasant emotional and physical discomfort. Such use in relief marks the most common occurrence of both behaviours and presents the highest risk situations, though this is somewhat less the case for college students than other demographic brackets. In contrast, heavy episodic drinking appears more likely than binge eating to take place in positive reinforcement contexts marked by pleasant

such literacy will be recognition of how interconnections among the diverse array of contributing factors have an impact on well-being for individuals and communities (Freedman et al., 2009).

The social-ecological model of public health and health promotion reflects the multidirectional complexity of dynamic interplay among factors operating within and across different levels from macro (societal) through micro (individual), so that environments affect people personally and corporately,

drinking behaviour as is a popular cultural ethos around alcohol in which it is viewed as an indispensible means for having fun with others and establishing gratifying social networks. Efforts to improve the ecosystem of the campus context itself will take different forms (DeJong & Langford, 2002; Toomey & Wagenaar, 2002; Toomey, Lenk & Wagenaar, 2007), including more regulatory policy positions (e.g., constraints on alcohol availability and promotion; cf. Wechsler & Nelson, 2008; Nelson et al., 2009, 2010) as well as attempts to cultivate and nurture a community of constructive, caring interaction (Ziemelis et al., 2002; Murphy et al., 2007; Borsari et al., 2007). Such an agenda will consider effective ways of providing alternative socialization and a variety of relationship-building settings, as well as engaging students in on-campus and community service opportunities (CARBC, 2010a).

Efforts that strengthen the relational fabric of a post-secondary school and enhance participation will be beneficial not only in helping to deter students from recourse to hazardous substance use, but also be something of a buffer against development of patterns of disordered eating. An approach similar to ï • ‡ƒŽ–Š”‘•‹–‰…Š‘ŽŽ would supply a more holistic rubric for encouraging positive attitudes a• †„‡Šƒ˜‹—”• ï‡ƒ¬ƒŽ‘•á trrrâƒ‰”á CARBC, 2010b; WHO, n.d.). Its thrust not only diffuses health education across the curriculum with orientation and training aids for teachers and staff, but also calls for examination of structures, policies and practices that work against rather than support health. It also urges strong collaborative working partnerships with other sectors of the community, including social and health services among others (e.g., community recreation and sport associations). Application of such a philosophy within the collegiate setting to create and sustain a wholesome ethos will employ like means of integrated education, community mobilization and administrative or organizational action to ensure positive interventions.

standards) that support or uphold a healthier outlook on personally and socially acceptable appearance

Attempts to increase media literacy and enhance body satisfaction can be usefully embedded within multi-component programs, universal or selected, to elevate self esteem, improve management skills and encourage positive relationship building (e.g., McVey et al., 2016). Prevention program reviews around disordered eating include some initiatives that aim to offer help on behavioural difficulties already occurring (as reflected in further discussion below under brief and more extended interventions), as well as efforts to reduce onset of future symptoms (e.g., Stice & Shaw,

suggests that some progress may be made simply by this build-up of self-esteem and appreciation of personal ability without necessarily needing to focus simultaneously on reducing particular symptoms, the strongest package would appear to be one that supplements or combines such a motivation enhancing thrust with tools that help clients work through strategies to implement change. On the one hand, attention might be paid to raising awareness of causes for concern, exploring ambivalence, assessing readiness, considering pros and cons, reflecting on values and dissonance in relation to those, recognizing and affirming capacity and competence, generating ideas for change and selecting from alternatives, and designing a plan. On the other side, a resource might clarify the pattern in question, introduce techniques that may be employed in a process of change, take up points of challenge and problem solving steps, as well as address relapse prevention matters, all with a view to imparting skills and raising literacy to access external helps (Cassin et al., 2008; Dunn et al., 2006).

Cognitive behavioural skills training

A stepped care approach in intervention responding to individual problems around substance use and/or disordered eating would first offer unsupervised self-help available in some form (e.g., referral to an online resource), proceed as required to more guided forms of such assistance in a group context, and move to private formats involving sessions with the person experiencing the difficulty. When the intervention ‹ • ‡ š – ‡ • † ‡ † „ ‡ › ‘ • † – Š ‡ ò „ ” ‹ ‡ ˆ ó ’ ƒ ” ƒ • ‡ – ‡ ” • ‘ ˆ ƒ • ‹ • few segments into a more prolonged series of engagements, with more intensive effort being placed on equipping clients, cognitive behavioural therapy (CBT) is the most widely attested treatment. On the eating disorders front, it essentially amounts to a continuation and amplification of the sorts of components alluded to above under skill development in brief intervention, ideally incorporating some of the motivational enhancement emphases in the process (Grilo et al., 2002; Sinha & ï ƒ Ž Ž ‡ › á t r r r â † ƒ • • ¬ ” ƒ ƒ • á t r r x â ƒ • • ‹ • ‡ – ƒ Ž ä á t r r z ä

For difficulties with substance use, particularly alcohol, the Alcohol Skills Training Program † ‡ ˜ ‡ Ž ‘ ’ ‡ † ƒ • „ › – Š ‡ • ‹ ˜ ‡ ” • ‹ – addictive Behaviors Research Center (Alan Marlatt and colleagues: Miller et al., 2001; Fromme et al., 1994; Kivlahan et al., 1990) is regarded as the CBT exemplar that has been most productively appropriated (Larimer & Cronce, 2007; Neighbors et al., 2006; Walters & Neighbors, 2005) and has inspired kindred programs (e.g., Fromme & Orrick, 2004). While there has been much less formulation and testing of initiatives aimed at reaching those experiencing difficulties with other drugs, indications are that the same key

However, though there can be direct overlap in dealing with factors that contribute to both substance and food use problems, complexity and diversity must be kept in view. Within eating disorder categories, there are different behaviours (e.g., restricting, bingeing, purging) that will reflect distinct considerations or interests on the part of various individuals and may link up with substance use in different ways. Use of stimulants can serve the cause of appetite suppression and weight reduction (cf. Piran & Robinson, 2006a,b; Herzog et al., 2006). Anticipated alcohol intake can take priority over and prompt lesser intake of food (in order to minimize aggregate calories), which circumstances can contribute to earlier intoxication and associated dangers as well as greater liability to internal physical damage. The context and effects of alcohol use can also limit or hinder sound choices on quality of food as well as compromise ability to regulate food intake and process it. Thus alcohol can abet both unhealthy dieting and excessive weight gain (Giles et al., 2009; Lloyd Richardson et al., 2008; Nelson et al., 2009). Binge eating tends to be a more private, solitary practice given the stigma attached to it, in contrast to the perceived acceptability of heavy drinking episodes in some public settings (Birch et al., 2007). Illegality of other substances will have some impact on their contexts of use. Exploration and discussion of predominant motives and incentives should be an integral component of brief intervention and more extended skills training for those struggling with such problems.

While CBT is widely accepted as a preferred therapy in regard to eating disorders, other forms of skill-enhancing care enjoy some advocacy too. Dialectical behaviour therapy (DBT) has been adapted from use with borderline personality disorder and shows some promise for dealing with regulation of negative affect and avoidant coping (Grilo et al., 2002; Telch, Agras & Linehan, 2001). DBT is one form among others of mindfulness training that is being explored for possible utility in intervention in both eating and drinking problems. The accent of mindfulness will be on non-Œ—†‰•‡•–ƒ Ž ƒ––‡•–ƒ•–emotional, cognitive and bodily experience in the present moment; training can foster utilization of this detached, decentred observational awareness and acceptance to

Reference Material

Adams, T.B. & Araas, T.E. (2006). Purging and alcohol-related effects in college women. International Journal of Eating Disorders, 39(3), 240-244.

Adlaf, E.M., Demers, A. & Gliksman, L. (Eds.) (2005). Canadian Campus Survey 2004. Toronto: Centre for Addiction and Mental Health. http://www.camh.net/research/population_life_course.html .

Anderson, D.A., Lundgren, J.D., Shapiro, J.R. & Paulosky, C.A. (2004). Assessment of eating disorders: Review and recommendations for clinical use. Behavior Modification, 28(6), 763-782.

Anderson, D.A., Martens, M.P. & Cimini, M.D. (2005). Do female college students who purge report greater alcohol use and negative alcohol-related consequences? International Journal of Eating Disorders, 37(1), 65-68.

Anderson, D.A., Simmons, A.M., Martens, M.P., Ferrier, A.G. & Sheehy, M.J. (2006). The relationship between disordered eating behavior and drinking motives in college-age women. Eating Behaviors, 7(4), 419-422.

Andreasen, A.R. (2004). A social marketing approach to changing mental health practices directed at youth and adolescents. Health Marketing Quarterly, 21(4), 51-75.

Anstine, D. & Grinenko, D. (2000). Rapid screening for disordered eating in college-aged females. Journal of Adolescent Health, 26(5), 338-342.

Baer, R.A. (2003). Mindfulness training as a clinical intervention: A conceptual and empirical review. Clinical Psychology: Science and Practice, 10(2), 125-143.

Bashford, J., Flett, R. & Copeland, J. (2010). The Cannabis Use Problems Identification Test (CUPIT): development, reliability, concurrent and predictive validity among adolescents and adults. Addiction, 105(4), 615-625.

Bauer, G., Davies, J.K., Pelikan, J., Noack, H., Broesskamp, U. & Hill, C. (2003). Advancing a theoretical model for public health and health promotion indicator development. European Journal of Public Health, 13(3), 107-113.

Becker, A.E., Franko, D.J., Nussbaum, K. & Herzog, D.B. (2004). Secondary prevention for eating disorders: The impact of education, screening and referral in a college-based screening program. International Journal of Eating Disorders, 36(2), 157-162.

Becker, A.E., Franko, D.L., Speck, A. & Herzog, D.B. (2003). Ethnicity and differential access to care for eating disorder symptoms. International Journal of Eating Disorders, 33(2), 205-212.

Benjamin, L. & Wulfert, E. (2005). Dispositional correlates of addictive behaviors in college women: Binge eating and heavy drinking. Eating Behaviors, 6(3), 197-209.

Beseler, C.L., Taylor, L.A. & Leeman, R.F. (2010). An item-response theory analysis of DSM-IV alcohol-use †‹•‘”†‡” …”‹–‡”‹ƒ ƒ•• ‰ó †”‹•Journal of Studies on Alcohol and Drugs, 71(3), 418-423.

Birch, C.D., Stewart, S.H. & Brown, C.G. (2007). Exploring differential patterns of situational risk for binge eating and heavy drinking. Addictive Behaviors, 32(3), 433-448.

Borsari, B., Murphy, J.G. & Barnett, N.P. (2007). Predictors of alcohol use during the first year of college: Implications for prevention. Addictive Behaviors, 32(10), 2062-2086.

Brennan, J. (2005). A selective prevention study: Decreasing body dissatisfaction and eating disorder symptomatology in sorority women using psychoeducation, social norms and social marketing strategies. http://etd.ohiolink.edu/send-pdf.cgi/Brennan%20Julie.pdf?osu1121286117

Bulik, C.M., Klump, K.L., Thornton, L., Kaplan, A.G <</Md

Fleming, M.F., Barry, K.L. & MacDonald, R. (1991). The Alcohol Use Disorders Identification Test (AUDIT) in a college sample. International Journal of the Addictions/Substance Use and Misuse, 26(11), 1173-1185.

Franko, D.L., Dorer, D.J., Keel, P.K., Jackson, S., Manzo, M.P. & Herzog, D.B. (2005). How do eating disorders and alcohol use disorder influence each other? International Journal of Eating Disorders, 38(5), 200-207.

Franko, D.L., Mintz, L.B., Villapiano, M., Green, T.C., Mainelli, D., Folensbee, L. et al. (2005). Food, mood and attitude: Reducing risk for eating disorders in college women. Health Psychology, 24(6), 567-578.

Freedman, D., Bess, K., Tucker, H., Boyd, D., Tuchman, A., & Wallston, K. (2009). Public health literacy defined. American Journal of Preventive Medicine, 36(5), 446-51.

Fromme, K. & Orrick, D. (2004). The lifestyle management class: A harm reduction approach to college drinking. Addiction Research and Theory, 12(4), 335-351.

Fromme, K., Marlatt, G.A., Baer, J.S. & Kivlahan, D.R. (1994). The alcohol skills training program: A group intervention for young adult drinkers. Journal of Substance Abuse Treatment, 11(2), 143-154.

Gadalla, T. & Piran, N. (2007a). Co-occurrence of eating disorders and alcohol use disorders in women: a meta analysis. ” … Š ‹ ˜ ‡ • ‘ ˆ ‘ • ‡ • ï • ‡ , 1 0 (4) , 1 3 3 - 1 4 0 . – Š

Gadalla, T. & Piran, N. (2007b). Eating disorders and substance abuse in Canadian men and women: a national study. Eating Disorders, 15(3), 189-203.

Garner, D.M., Olmsted, M.P., Bohr, Y. & Garfinkel, P.E. (1982). The eating attitudes test: Psychometric features and clinical correlates. Psychological Medicine, 12(4), 871-878.

Gilchrist, G., Gruer, L. & Atkinson, J. (2007). Predictors of neurotic symptom severity among female drug users in Glasgow, Scotland. Drugs: Education, Prevention and Policy, 14(4), 347-365.

Giles, S.M., Champion, H., Sutfin, E.L., McCoy, T.P. & Wagoner, K. (2009). Calorie restriction on drinking days: an examination of drinking consequences among college students. Journal of American College Health, 57(6), 603-609.

Grant, B.F., Dawson, D.A., Stinson, F.S., Chou, S.P., Dufour, M.C., Compton, W. & Pickering, R.P. (2004). The 12-month prevalence and trends in DSM-

c u6 v el0 05pÍ 1 ET abu Ünking coh%nOA.erV cC{N±O F48Î r Cp ' Vñ%Þ …Á:C²"F V†•6a:C pcÁ` ¤3A ' “£dpt ¤:Õnëñ Vñ 5A:CH

Kivlahan, D.R., Marlatt, G.A., Fromme, K., Coppel, D.B., & Williams, E. (1990). Secondary prevention with college drinkers: Evaluation of an alcohol skills training program. Journal of Consulting and Clinical

McMillen, B.A., Hillis á ä ä ¬ ”‘™•á ä ä trr{ ä ‘ŽŽ‡‰‡ •–—†‡•–•ï ”‡•’‘••‡• –‘ ƒ and maximum blood alcohol concentration calculated from a timeline followback questionnaire. Journal of Studies on Alcohol and Drugs, 70(4), 601-605.

McVey, G., Kirsh, G., Maker, D., Walker, K.S., Mullane, J., Laliberte, M. et al. (2010). Promoting positive body image among university students: A collaborative pilot study. Body Image, 7(3), 200-204.

Measelle, J.R., Stice, E. & Hogansen, J.M. (2006). Developmental trajectories of co-occurring depressive, eating, antisocial, and substance abuse problems in female adolescents. Journal of Abnormal Psychology, 115(3), 524-538.

Miller, E., Kilmer, J.R., Kim, E.L., Weingardt, K.R. & Marlatt, G.A. (2001). Alcohol skills training for college •–—†‡•–•ä • ‘•–‹á ä äá ‘Ž„›á ä Adolescents, Alcohol and Substance Abuse: Reaching Teens Through Brief Interventions (pp. 183-215). New York: Guilford Press.

Miller, W.R. & Rollnick, S. (2009). Ten things that motivational interviewing is not. Behavioural and Cognitive Psychotherapy, 37(2), 129-140.

Miller, W.R. & Rollnick, S. (2004). Talking oneself into change: motivational interviewing, stages of change, and therapeutic process. Journal of Cognitive Psychotherapy: An International Quarterly, 18(4), 299-308.

Miller, W.R. & Rollnick, S. (2002). Motivational interviewing: Preparing people to change. New York: Guilford Press, 2nd ed.

Murphy, J.G., Correia, C.J. & Barnett, N.P. (2007). Behavioral economic approaches to reduce college student drinking. Addictive Behaviors, 32(11), 2573-2585.

National Institute on Alcohol Abuse and Alcoholism (NIAAA, 2004). NIAAA Council Approves Definition of Binge Drinking. NIAAA Newsletter Winter 2004, number 3, p.3. NIH Publication No. 04-5346. Available at http://pubs.niaaa.nih.gov/publications/Newsletter/winter2004/Newsletter_Nu mber3.pdf.

Neighbors, C., Larimer, M.E., Lostutter, T.W. & Woods, B.A. (2006). Harm reduction and individually focused alcohol prevention. International Journal of Drug Policy, 17(4), 304-309.

Nelson, M.C., Lust, K., Story, M. & Ehlinger, E. (2009). Alcohol use, eating patterns, and weight behaviors in a university population. American Journal of Health Behavior, 33(3), 227-237.

Nelson, T.F., Toomey, T.L., Lenk, K.M., Erickson, D.J. & Winters, K.C. (2010). Implementation of NIAAA College Drinking Task Force recommendations: How are colleges doing 6 years later? Alcoholism: Clinical & Experimental Research, 34(10), 1687-1693.

Nelson, T.F., Xuan, Z., Lee, H., Weitzman, E.W. & Wechsler, H. (2009). Persistence of heavy drinking and ensuing consequences at heavy drinking colleges. Journal of Studies on Alcohol and Drugs, 70(5), 726-734.

NIAAA (2007). What Colleges Need to Know Now: An Update on College Drinking Research http://www.collegedrinkingprevention.gov/1CollegeBulletin-508_361C4E.pdf

NIAAA (2002a). A Call to Action: Changing the Culture of Drinking at U.S. Colleges http://www.collegedrinkingprevention.gov/NIAAACollegeMaterials/TaskForce/TaskForce_TOC.aspx. http://www.collegedrinkingprevention.gov/media/TaskForceReport.pdf

NIAAA (2002b). High-risk drinking in college: What we know and what we need to learn. Final report of the panel on contexts and consequences. http://www.collegedrinkingprevention.gov/media/FINALPanel1.pdf

NIAAA (2002c). How to reduce college drinking: Use proven strategies, fill research gaps. Final report of the panel on prevention and treatment. http://www.collegedrinkingprevention.gov/media/FINALPanel2.pdf

Nutbeam, D. (2008). The evolving concept of health literacy. Social Science & Medicine, 67(12), 2072-78.

ï ”‹‡•á ä äá … ‘›á ä äá Šƒ•’‹‘•á äá ‹–”ƒá äá ‘„„‹••á äá ‡—•…ŠŽ• question about drunkenness to detect college students at risk for injury. Academic Emergency Medicine, 13(6), 629-636.

ï ‡ƒ Á. & Abraham, S. (2002). Eating and exercise disorders in young college men.

Rootman, I., Goodstadt, M., Potvin, L. & Springett, T. (2001). A framework for health promotion evaluation. In I. Rootman, M. Goodstadt, B. Hyndman et al. (eds.), Evaluation in health promotion: Principles and perspectives (pp. 7-38). Copenhagen: World Health Organization.

Rush, C.C., Becker, S.J. & Curry, J.F. (2009). Personality factors and styles among college students who binge eat and drink. Psychology of Addictive Behavior, 23(1), 140-145.

Sansone, R.A., Fine, M.A. & Nunn, J.L. (1994). A comparison of borderline personality symptomatology and self-destructive behavior in women with eating, substance abuse, and both eating and substance abuse disorders. Journal of Personality Disorders, 8(3), 219-228.